

# Air pollution positively correlates with daily stroke admission and in hospital mortality: a study in the urban area of Como, Italy

Simone Vidale · A. Bonanomi · M. Guidotti ·  
M. Arnaboldi · R. Sterzi

Received: 28 October 2008 / Accepted: 14 December 2009 / Published online: 30 January 2010  
© Springer-Verlag 2010

**Abstract** Some current evidences suggest that stroke incidence and mortality may be higher in elevated air pollution areas. Our study examined the hypothesis of a correlation between air pollution level and ischemic stroke admission and in Hospital mortality in an urban population. Data on a total of 759 stroke admissions and 180 deaths have been obtained over a 4-year period (2000–2003). Five air ambient particles have been studied. A general additive model estimating Poisson distribution has been used, adding meteorological variables as covariates. NO<sub>2</sub> and PM<sub>10</sub> were significantly associated with admission and mortality ( $P$  value < 0.05) and with estimated RR of 1.039 (95% CI 1.066–1.013) and 1.078 (95% CI 1.104–1.052) for hospital admission at 2- and 4-day lags, respectively. In conclusion, this study suggests an association between short-term outdoor air pollution exposure and ischemic stroke admission and mortality.

**Keywords** Ischemic stroke · Risk factor · Air pollution

---

S. Vidale (✉) · M. Arnaboldi  
Neurology Department, Sant'Anna Hospital,  
Via Napoleona 60, 22100 Como, Italy  
e-mail: simone.vidale@hsacomo.org; simon.vidale@gmail.com

A. Bonanomi  
Statistic Sciences Department,  
Cattolica Sacro Cuore University, Milan, Italy

M. Guidotti  
Neurology Department, Valduce Hospital, Como, Italy

R. Sterzi  
Neurology Department, Niguarda Ca' Granda Hospital,  
Milan, Italy

## Introduction

Epidemiological studies demonstrated that high level of outdoor air pollution can increase the incidence and the mortality risk for cardiovascular diseases [1, 2]. Based on these findings, some studies tried to demonstrate an association between elevated short-term outdoor air pollution and cerebrovascular diseases. The evidence suggests that stroke hospital admissions are higher in areas with elevated levels of outdoor air pollutants. This is due to the combined acute and chronic effects of the exposure to air pollution on stroke risk. There is an increasing interest in using hospital admission data and air pollutant short-term effects. Time series studies show a statistically significant positive association between daily levels of particulate matter and other gaseous pollutants and hospitalization for cerebrovascular events.

Studies conducted in Seoul, Korea, in Taiwan, and in Europe have shown a significant association between some air pollutants and stroke hospital admissions and mortality [3–7]. On the other hand, some works reported lack of association [8, 9].

We investigate the association between short-term outdoor air pollutant levels and ischemic stroke hospital admissions and mortality in Como urban area citizens, over a 4-year period (2000–2003) and before the setting of the two Stroke Units now existing in the town.

## Materials and methods

Como is situated in the North-West of Italy and its population is about 85,000 people. Como downtown is near the lake and it is surrounded by not very high mountains causing a poor ambient-air circulation and persistent high

level of air pollutants. Data about hospital admissions for ischemic stroke and mortality-related are obtained from the Stroke Registry of the two principal City Hospitals. To identify some other cases, we analyzed computerized records of daily clinic hospital admissions, using the codes 433 and 434 of the International Classification of Disease, 9th revision (ICD-9). For these cases we have no data about vascular risk factors and stroke severity. Two air-quality monitoring stations were established in Como and controlled by the Regional Agency for the Ambient Protection (ARPA). These monitoring stations are automated and can provide daily reading and saving of SO<sub>2</sub>, NO<sub>2</sub>, CO, O<sub>3</sub> and PM<sub>10</sub>. We obtained daily data about ambient temperature, atmospheric pressure, wind speed, and precipitation level from two meteorological stations.

The associations between daily stroke admissions and mortality in Como and air pollutions, as well as weather variables, were analyzed with a general additive model (GAM) estimating Poisson distribution (the dependent variable was rare and was assumed to be Poisson distributed) [10]. First, GAM was used to study associations between stroke admission and air pollution covariates (SO<sub>2</sub>, NO<sub>2</sub>, CO, O<sub>3</sub> and PM<sub>10</sub>) with different time-delay (1–5 days). Second, meteorological covariates were introduced in the model with different time-delay. The final model was determined using a backward selection process in which model covariates were examined for their statistical significance and no interactions were considered in order to avoid estimating too many parameters.

The odds ratio and the estimate relative risk for admission were calculated for each air pollutant considering exposed days and not-exposed days and the number of days with registered events [11]. In the analysis we considered alarm levels of each pollutant as reported in the actual European laws. Using a  $\chi^2$  test, a different lag time between 0 and 5 days was considered.

## Results

There were a total of 759 ischemic stroke admissions and 180 deaths during hospitalization. On average, there were 2.2 ischemic strokes every 1,000 Como citizens/year, with 23.6% mortality year-rate. Mean age was  $68 \pm 13$  years, with a prevalence of male (56.7%). Analyzing air pollutant levels, a cyclical trend was found, with higher levels during autumn and winter, except for the ozone which showed the pick in the summer time.


Analyzing hospital stroke admissions and each data, higher ischemic event rate was found in cool days than in the warm ones. The statistical model showed an inverse association ( $P < 0.05$ ).

Using the statistical model, we observed a significant association between ischemic stroke admissions and the trend of values of NO<sub>2</sub>, CO, and O<sub>3</sub> ( $P < 0.001$ ). After meteorological data inclusion in the statistical model, among these pollutants, NO<sub>2</sub> remained the only agent with a positive association ( $P < 0.05$ ). Besides that PM<sub>10</sub> was associated with ischemic stroke hospital admissions ( $P < 0.05$ ).

Using the same statistical model, a positive association was observed between daily mortality and NO<sub>2</sub> level ( $P < 0.01$ ) with 1-day delay after pollutant exposure and with PM<sub>10</sub> levels ( $P < 0.05$ ) with 5-day delay. No other air pollutant was associated with daily mortality. Contour analysis of PM<sub>10</sub> and NO<sub>2</sub> with daily stroke admission with 1-day lag was showed in the generalized additive model plot (Fig. 1).

We observed estimated relative risks of 1.057 (95% CI 1.083–1.031) and 1.039 (95% CI 1.065–1.013) for hospital admission for PM<sub>10</sub> and SO<sub>2</sub> concentrations, respectively, on the same day. We found the maximum estimated relative risk of 1.078 (95% CI 1.104–1.052) for PM<sub>10</sub> with a 4-day lag. We observed statistically significant association between hospital ischemic stroke admission and NO<sub>2</sub> with a 2-day lag (RR 1.039, 95% CI 1.066–1.013). At the same time lag, we found the best estimated relative risk for SO<sub>2</sub> with 1.058 (95% CI 1.085–1.032).

For CO, we observed a significant relative risk only with 1-day lag (RR 1.060, 95% CI 1.087–1.034).



**Fig. 1** Contour analysis of PM<sub>10</sub> and NO<sub>2</sub> with daily stroke admission at different lag days

**Table 1** Estimated relative risk and 95% confidence interval of hospital admission for ischemic stroke in exposed and not-exposed days in PM<sub>10</sub>, NO<sub>2</sub>, SO<sub>2</sub>, CO and O<sub>3</sub> in single-pollutant model

Lag	PM <sub>10</sub>		NO <sub>2</sub>		SO <sub>2</sub>		CO		O <sub>3</sub>	
	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI	RR	95% CI
0	1.057	1.083–1.031	0.990	1.015–0.963	1.039	1.065–1.013	0.976	1.003–0.950	0.920	0.947–0.894
1	1.010	1.036–0.98	1.010	1.035–0.983	1.031	1.057–1.004	1.060	1.087–1.034	0.881	0.907–0.855
2	0.990	1.016–0.96	1.039	1.066–1.013	1.058	1.085–1.032	0.969	0.995–0.942	0.936	0.962–0.910
3	1.044	1.07–0.98	0.981	1.007–0.955	1.041	1.068–1.015	1.005	1.031–0.979	0.870	0.896–0.844
4	1.078	1.10–1.052	0.985	1.012–0.959	1.042	1.069–1.016	1.021	1.048–0.995	0.911	0.937–0.887
5	1.000	1.02–0.96	1.017	1.043–0.991	1.017	1.044–0.991	1.005	1.032–0.979	0.867	0.893–0.841

No significant estimated relative risk was observed for O<sub>3</sub> (Table 1).

## Discussion

To the best of our knowledge, this is the first Italian study to report a specific association between daily ischemic stroke hospital admissions and mortality, and air pollution. In fact, the previous meta-analysis of Italian studies was published without specific references to cerebrovascular disease [12].

As reported in several studies conducted in other countries, in the present work, a positive association between NO<sub>2</sub> and PM<sub>10</sub> levels and ischemic stroke hospital admissions and mortality was observed. As reported in the analysis of the two air pollutants (PM<sub>10</sub> and NO<sub>2</sub>), the 2-day lag seems important to evaluate the short-term accumulated dose of air pollutants and to avoid other confounding factors in the hospital admission. It is interesting to note that meteorological data are very important for a correct analysis of the association and the most important variable seems to be the temperature. Differently from published studies in the South-East of Asia, we observed a negative association with the temperature because the number of stroke hospital admissions was higher during cool months, in particular, December, January and February. This evidence is not described by other similar studies in European or North-American countries [5–7]. In the urban area of Como, it is possible to observe higher levels of air pollutants during this period because of the turning on heat and the increasing car traffic. The increase in relative risk of hospital admission for ischemic stroke with air pollutant rising is in the order of 3–5%.

In the past, there have been many epidemiological and toxicological studies concerned with the potential mechanism linking air pollution and cardiovascular diseases, including stroke. Frampton et al. [13] hypothesized a

systemic inflammation and vasoconstriction with an expression of leukocytes, endothelial adhesion molecules, oxidants and interleukins, induced by changes in vascular function due by particle acidity or particles with transition metals or ultrafine particles.

As other published studies, this work has some limitations. First, there is a potential selection bias due to clinically unrecognized stroke cases or stroke treated in other hospitals. Second, epidemic diseases were not included in the statistical model because data were not available. However, we observed an increased level of hospital stroke admissions before the classical beginning of epidemic influenza or getting worse again of other upper and lower respiratory diseases. Some recent studies have suggested the role of acute infections as trigger factor for cerebrovascular accident, but the relationship is not clear. However, other studies are actually ongoing to search a direct association between cerebral ischemic event and inflammation related to infections. Third, there are ecological biases which cannot be ruled out. Fourth, individual exposure was assigned as air pollutant level from fixed outdoor monitoring stations. Measurement errors resulting from differences between population-average exposure and ambient levels cannot be avoided. However, this kind of measurement error is known to cause a bias toward null and underestimates the pollution effects (Berkson type error) [4]. Finally, we have no data available on hypertension prevalence or other stroke risk factors that could potentially have mislead the associations observed. However, many individual vascular risk factors do not vary significantly over time or vary very slowly and they are unlikely to be confounding factors in time series study [4].

The present study shows the positive association between some air pollutants and ischemic stroke. Although this population is not large, these results suggest that air pollution could be a potentially modifiable risk factor for stroke. By focusing on the targeting policy interventions at high pollution areas, we could reduce stroke-related health problems more significantly.

## References

1. Brook RD, Franklin B, Cascio W, Hong Y, Howard G, Lipsett M, Luepker R, Mittleman M, Samet J, Smith SC Jr, Tager I (2004) Expert Panel on Population and Prevention Science of the American Heart Association. Air pollution and cardiovascular disease: a statement for healthcare professionals from the Expert Panel on population and prevention science of the American Heart Association. *Circulation* 109:2655–2671
2. Miller KA, Siscovick DS, Sheppard L, Shepherd K, Sullivan JH, Anderson GL, Kaufman JD (2007) Long-term exposure to air pollution and incidence of cardiovascular events in women. *New Eng J Med* 356:447–458
3. Tsai SS, Goggins WB, Chiu HF, Yang CY (2003) Evidence for an association between air pollution and daily stroke admissions in Kaohsiung, Taiwan. *Stroke* 34:2612–2616
4. Hong YC, Lee JT, Kim LH, Kwon HJ (2002) Air pollution. A new risk factor in ischemic stroke mortality. *Stroke* 33:2165–2169
5. Maheswaran R, Haining RP, Brindley P, Law J, Pearson T, Fryers PR, Wise S, Campbell MJ (2005) Outdoor air pollution and stroke in Sheffield, United Kingdom. A small-area level geographical study. *Stroke* 36:239–243
6. Hoek G, Brunekreef B, Goldbohm S, Fisher P, van den Brandt PA (2003) Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort study. *Lancet* 360:1203–1209
7. Piver WT, Ando M, Ye F, Portier CJ (1999) Temperature and air pollution as risk factors for heat stroke in Tokyo, July and August 1980–1995. *Environ Health Perspect* 107:911–916
8. Poloniecki JD, Atkinson RW, de Leon AP, Anderson HR (1997) Daily time series for cardiovascular hospital admissions and previous day's air pollution in London, UK. *Occup Environ Med* 54:535–540
9. Le Tertre A, Medina S, Samoli E, Forsberg B, Mechelozzi P, Boumghar A, Vonk JM, Bellini A, Atkinson R, Ayres JG, Sunyer J, Schwartz J, Katsouyanni K (2002) Short term effects of particulate air pollution on cardiovascular diseases in eight European cities. *J Epidemiol Community Health* 56:773–779
10. Zhang J, Yu KF (1998) What's the relative risk? A method of correcting the odds ratio in cohort studies of common outcomes. *JAMA* 280:1690–1691
11. Biggeri A, Bellini P, Terracini B (2001) Meta-analysis of the Italian studies on short-term effects of air pollution. *Epidem Prevent Suppl* 2:2–71
12. Wellenius GA, Schwartz J, Mittleman MA (2005) Air pollution and hospital admissions for ischemic and hemorrhagic stroke among medicare beneficiaries. *Stroke* 36:2549–2553
13. Frampton MW, Stewart JC, Oberdörster G, Morrow PE, Chalupa D, Pietropaoli AP, Frasier LM, Speers DM, Cox C, Huang LS, Utell MJ (2006) Inhalation of ultrafine particles alters blood leukocyte expression of adhesion molecules in humans. *Environ Health Perspect* 114(1):51–58